



NASA Langley Research Center

Trajectory Reconstruction for the Genesis Entry

Prasun Desai

Garry Qualls

Mark Schoenenberger

NASA Langley Research Center


June 28, 2005



Genesis Mission Background

NASA Langley Research Center

- Fifth Discovery mission
 - Launched on August 8, 2001
 - Inserted in to a halo orbit about the sun-Earth libration point (L1)
 - Collected solar wind particles for ~29 months
 - First mission to return samples from beyond the Earth-moon system
- Maneuver and targeting procedures placed capsule on desired entry flight-path on morning of September 8, 2004 at 15:52:47 UTC






Genesis Entry Live!

NASA Langley Research Center

Capsule mass = 205.6 kg





Genesis Entry Live!

NASA Langley Research Center

QuickTime™ and a
decompressor
are needed to see this picture.




Genesis Nominal Entry Sequence

NASA Langley Research Center

Entry $V_I = 11.04$ km/s

Entry $\gamma_I = -8.0$ deg






Genesis Impact Location

NASA Langley Research Center

Target Location: (246.4667 deg, 40.2 deg)

Nominal Landing Location: (246.4819 deg, 40.2092 deg)

Final Impact Location: (246.4919 deg, 40.1278 deg)





Genesis Reconstruction Effort

NASA Langley Research Center


- Genesis capsule landed very close to the desired target
 - Understanding hypersonic flight performance is of great interest
 - Specifically, assessing how the pre-entry predictions of flight dynamics, aerodynamics, and aerothermodynamics
 - Identify Mach number for onset of capsule tumble
- Only limited data exists to perform reconstruction
 - No onboard sensors on capsule
 - Only available data source is from UTTR radar tracking stations
 - Video and tracking data
 - Balloon measurement of atmospheric properties up to 34 km four hours prior to entry



Density Comparison to GRAM Model

NASA Langley Research Center

- Global Reference Atmosphere Model (GRAM) used for pre-entry prediction






Density Comparison to GRAM Model (cont'd)

NASA Langley Research Center

- Variation of $\pm 2.5\%$ observed on entry day
- Corresponds to $\sim 1.5\text{-}\sigma$ profile from GRAM variations






Wind Comparison to GRAM Model

NASA Langley Research Center

- Variation of ± 5 m/sec observed on entry day






Wind Comparison to GRAM Model (cont'd)

NASA Langley Research Center


- Sustained wind to the East observed (~ 27 m/sec at 12 km)
- Corresponds to $\sim 1.5\text{-}\sigma$ profile from GRAM variations





Capsule Luminance from Infrared Video


NASA Langley Research Center





Frequency Contours of Luminance Data

NASA Langley Research Center






Capsule Luminance from Infrared Video

NASA Langley Research Center

- Onset of tumble at Mach 0.9






Pre-Entry Simulation Capsule Attitude Prediction

NASA Langley Research Center

- Results from video analysis and use of trajectory simulation help corroborate capsule aerodynamics in supersonic regime (Mach 2.2 - 1.0)






Capsule Deceleration Comparison (Supersonic Regime)

NASA Langley Research Center

- Timelines of simulation and tracking data aligned at Mach 1
- Good agreement indicates aerodynamic database accurately captures capsule drag






Capsule Frequency Comparison (Supersonic Regime)

NASA Langley Research Center

- Dominant frequency correlate well over range where data is available





Capsule Static Stability Assessment (Supersonic Regime)

NASA Langley Research Center

- With confidence in oscillation frequency, an assessment of capsule static stability can be made since

$$\text{Freq} \propto \text{dynamic pressure} * (\text{pitching moment slope})^{0.5}$$

- Appears to be good agreement between measured data and pre-entry predicted dynamics pressure variation
 - Assertion depends on agreement between predicted drag and measured drag and agreement of density profile
 - Accuracy of drag shown by comparison of deceleration profile
 - Balloon measured density profile shown to be within $\pm 2.5\%$ of pre-entry profile
- Consequently, agreement of frequencies indicates aerodynamics database reasonably predicted static stability



Hypersonic Capsule Attitude Assessment

NASA Langley Research Center

- Since there was no onboard sensor data, capsule hypersonic attitude cannot be determined
 - Attitude must be inferred from observation of recovered heatshield
- There is very little, if any, charring of the shoulder region or aftbody of TPS
 - Inspection of forebody shows charring patterns that imply symmetry heating
 - Observations suggest that attitude must have been only a few degrees
 - Angle-of-attack is estimated to be no larger than $2.1^{\circ} \pm 1.4^{\circ}$
 - Pre-entry simulation predicted angle-of-attack of 1.38° with a maximum of 3°
- Overall observations support the assertion that the aerodynamics database reasonably predicted hypersonic static stability



Capsule Shoulder and Aftbody Region

NASA Langley Research Center





Capsule Forebody Heatshield


NASA Langley Research Center





Phosphor Thermography Tunnel Tests

NASA Langley Research Center





Trajectory Reconstruction

NASA Langley Research Center

- Since there was no onboard sensor data, a “traditional” trajectory reconstruction could not be performed
 - Therefore, a Best Estimated Trajectory (BET) is calculated
- The BET is split into two phases: hypersonic and subsonic flight
- For hypersonic flight, only two data sources were available
 - Final Navigation entry state at atmospheric interface
 - Latitude and longitude data from UTTR radar tracking stations at the pre-entry predicted drogue deployment time
 - UTTR radar tracking stations acquired capsule from 34 km through impact
- With confidence in these two endpoints, a hypersonic trajectory is calculated using the pre-entry simulation



Trajectory Reconstruction (cont'd)

NASA Langley Research Center

- Within the trajectory simulation, a multiplier on capsule drag was applied as the control parameter to determine the variation needed to patch the two endpoint conditions
 - An drag reduction of 8.1% is calculated
 - Altitude is within 380 m between simulation and tracking data
- The 8.1% reduction in drag can arise form multiple sources
 - Mis-prediction in entry state, capsule C_D , or atmospheric density
 - Final entry state was confirmed to be extremely accurate by STRATCOM
 - No measure of density available above 34 km
 - So, relative contributions between density and C_D cannot be determined
- However, an estimate for the hypersonic density can be approximated if an uncertainty in capsule C_D is assumed



Trajectory Reconstruction (cont'd)


NASA Langley Research Center

- Typical C_D uncertainty in the hypersonic regime is ± 4 ($3\text{-}\sigma$)
 - If a capsule C_D is assumed to be 1.5% low (a $1\text{-}\sigma$ occurrence), an estimate of atmospheric density above 34 km can be calculated
- With such an assumption, the atmospheric density encountered during the hypersonic flight can be approximated to be 6.6% lower than the pre-entry profile from GRAM
 - This estimate corresponds to $1.5\text{-}\sigma$ low profile from the GRAM variations
- Process for subsonic trajectory estimate is described in the paper



Genesis Best Estimated Trajectory

NASA Langley Research Center





Summary

NASA Langley Research Center

- An overview of the reconstruction analyses is described
- Atmosphere (density and winds) encountered during the entry were approximately 1- σ profiles
- Analysis on infrared video footage estimated the onset of capsule tumble at Mach 0.9
- Observations of the recovered heatshield indicated small attitude during hypersonic flight
- Overall assertion is that the Genesis entry flight performance was close to the nominal pre-entry predictions
- Consequently, the design principles and methodologies utilized for the flight dynamics, aerodynamics, and aerothermodynamics analyses were corroborated